

- KELLOGG, C. E. 1936. Development and Significance of the Great Soil Groups in the United States. U. S. Dept. Agr. Misc. Pub. 229.
- KITFIELD, A. 1974. A range extension for *Woodwardia areolata*. *Rhodora* 76:312.
- KOEVENIG, J. L. 1976. Effect of climate, soil, physiography and seed germination on the distribution of the River Birch (*Betula nigra*). *Rhodora* 78:420-437.
- KREBS, C. J. 1972. Ecology, the Experimental Analysis of Distribution and Abundance. Harper and Row, New York, NY.
- LITTLE, E. L. 1971. Atlas of United States Trees, vol. I. Conifers and Important Hardwoods. U. S. Dept. Agr. Misc. Pub. 1146.
- NATIONAL ATMOSPHERIC AND OCEANOGRAPHIC INSTITUTE. 1980. Climate of the States, 2 vols. Dept. of Commerce, Washington, DC.
- OGDEN, E. C., F. H. STEINMETZ, and F. HYLAND. 1948. Checklist of the Vascular Plants of Maine. *Bull. Josselyn Bot. Soc. Maine* 8.
- PIELOU, E. C. 1979. Biogeography. Wiley and Sons, New York, NY.
- SALISBURY, E. J. 1926. The geographical distribution of plants in relation to climatic factors. *Geogr. J.* 67:312-335.
- THORNTHWAITE, C. W. 1948. An approach to a rational classification of climate. *Geogr. Rev.* 38:55-94.
- WAGNER, W. H., Jr. 1963. Pteridophytes of the Mountain Lake area, Giles County, Virginia, including notes from Whitetop Mountain. *Castanea* 28:113-150.
- , and D. JOHNSON. 1981. Natural history of the Ebony Spleenwort, *Asplenium platyneuron* (Aspleniaceae) in the Great Lakes area. *Canad. Field Nat.* 95:156-166.
- WHERRY, E. T. 1921. The soil reactions of ferns of woods and swamps. *Amer. Fern J.* 11:5-16.
- . 1964. The Southern Fern Guide. Doubleday Nature Guide Series. Doubleday, Garden City, NY.

## REVIEW

**FERNS AND FERN ALLIES OF THE DRIFTLESS AREA OF ILLINOIS, IOWA, MINNESOTA AND WISCONSIN**, by James H. Peck. Milwaukee Public Museum Contributions in Biology and Geology 53:1-140. 1982. \$13.50 postpaid. —The driftless area, a pocket in the upper midwestern landscape which escaped at least the more recent glaciations, holds, by virtue of its topography and age of its habitats, a diverse and interesting array of plants. It has been the subject of botanical study for many years. The present account is by far the most complete and useful of any we have had for the pteridophytes, which number 73 species, 13 hybrids, and 6 infraspecific taxa. Peck's treatment includes a useful introduction, a discussion of the affinities of the flora with special reference to the interesting disjunct *Thelypteris simulata*, a systematic list with synonyms, statement of habitat, and specimen citations, a key to the genera and species of pteridophytes found in the driftless area, and a large and useful section of literature cited. Ninety pages are devoted to full-page county distribution maps for the driftless area ferns that include all of Minnesota, Wisconsin, Iowa, Illinois, and Missouri. The maps not only place the ferns of the driftless area in the context of the surrounding region, but they are sure to stimulate searches for pteridophytes in counties where the first records are yet to be obtained.—D.B.L.